

Spaces of Morphisms From a Projective Space to a Toric Variety

Espacios de morfismos de un espacio proyectivo a una variedad tórica

JACOB MOSTOVOY^{1,✉}, ERÉNDIRA MUNGUÍA-VILLANUEVA²

¹CINVESTAV-IPN, México, D.F., México

²Institut de Mathématiques de Jussieu, Paris, France

ABSTRACT. In this note we study the space of morphisms from a complex projective space to a compact smooth toric variety X . It is shown that the first author's stability theorem for the spaces of rational maps from $\mathbb{C}\mathbb{P}^m$ to $\mathbb{C}\mathbb{P}^n$ extends to the spaces of continuous morphisms from $\mathbb{C}\mathbb{P}^m$ to X , essentially, with the same proof. In the case of curves, our result improves the known bounds for the stabilization dimension.

Key words and phrases. Toric variety, Stone-Weierstrass Theorem, Spaces of toric morphisms, simplicial resolution.

2010 Mathematics Subject Classification. 58D15, 32Q55.

RESUMEN. En esta nota se estudia el espacio de morfismos de un espacio proyectivo complejo a una variedad tórica compacta no singular X . Se prueba que el teorema de estabilidad, demostrado por el primer autor para los espacios de funciones racionales de $\mathbb{C}\mathbb{P}^m$ a $\mathbb{C}\mathbb{P}^n$, se extiende a los espacios de morfismos continuos de $\mathbb{C}\mathbb{P}^m$ a X , esencialmente con la misma demostración. En el caso de las curvas, nuestro resultado mejora las cotas conocidas para la dimensión de la estabilización.

Palabras y frases clave. Variedad tórica, espacios de morfismos tóricos, Teorema de Stone-Weierstrass, resolución simplicial.

1. Introduction

Given two complex algebraic varieties one can ask how well the space of continuous algebraic morphisms between them approximates the space of all corresponding continuous maps. This question was posed by G. Segal in [20] where he

studied the spaces of holomorphic maps from a Riemann surface to a complex projective space. He proved the following result. Let S be a Riemann surface of genus g , $F_d(S, \mathbb{C}\mathbb{P}^n)$ the space of rational functions of degree d from S to $\mathbb{C}\mathbb{P}^n$, and $M_d(S, \mathbb{C}\mathbb{P}^n)$ the corresponding space of continuous maps. Similarly, denote by $F_d^*(S, \mathbb{C}\mathbb{P}^n)$ and $M_d^*(S, \mathbb{C}\mathbb{P}^n)$ the respective spaces of basepoint-preserving maps.

Theorem 1.1 ([20]). *The inclusions*

$$F_d(S, \mathbb{C}\mathbb{P}^n) \rightarrow M_d(S, \mathbb{C}\mathbb{P}^n) \quad \text{and} \quad F_d^*(S, \mathbb{C}\mathbb{P}^n) \rightarrow M_d^*(S, \mathbb{C}\mathbb{P}^n)$$

are homology equivalences up to dimension $(d-2g)(2n-1)$ for all $g \geq 0$. When $g = 0$ these maps are also homotopy equivalences up to the same dimension.

Segal conjectured that his results could be generalized much further to include spaces of curves in algebraic varieties more general than projective spaces. Such generalizations were obtained in subsequent works by many authors: Kirwan [14], Gravesen [9], Guest [10, 11], Mann and Milgram [15, 16], Hurtubise [13], Boyer, Hurtubise and Milgram [2], Cohen, Lupercio and Segal [4], and others. At the moment, the strongest results for the case when $g = 0$ (rational curves) and the target is a finite-dimensional variety X are those of Guest [11] for the case when X is a (possibly singular) toric variety and that of Boyer, Hurtubise and Milgram [2] for the case when X is a smooth Kähler manifold with a holomorphic action of a connected complex solvable Lie group which has an open dense orbit on which the group acts freely. A very general conjecture on this subject is given in the paper of Cohen, Jones and Segal [3]. Several papers [12, 17] give real versions of Segal's theorem (the first real version being due to Segal himself in [20]).

Segal's theorem can also be generalized to the case when the domain of the maps has dimension greater than 1. Using Vassiliev's simplicial resolutions, the first author in [18] and [19] proved an analog of Segal's theorem for the spaces of continuous rational maps from $\mathbb{C}\mathbb{P}^m$ to $\mathbb{C}\mathbb{P}^n$ for $1 \leq m \leq n$ (see also a real version in [1]).

The purpose of this note is to show that the proof given in [18, 19] extends to the spaces of maps from $\mathbb{C}\mathbb{P}^m$ to a smooth compact toric variety. We shall use the homogeneous coordinates for a toric variety X and the description of the morphisms $\mathbb{C}\mathbb{P}^m \rightarrow X$ given by Cox in [5] and [6]: a morphism $\mathbb{C}\mathbb{P}^m \rightarrow X$ is given in these coordinates by a collection of homogeneous polynomials with certain properties. Let P_f be the space of morphisms from $\mathbb{C}\mathbb{P}^m$ to X given by polynomials of fixed degrees, and whose restriction to a fixed hyperplane $\mathbb{C}\mathbb{P}^{m-1}$ coincides with a given morphism $f : \mathbb{C}\mathbb{P}^{m-1} \rightarrow X$. Consider the space of all continuous maps from $\mathbb{C}\mathbb{P}^m$ to X whose restriction to $\mathbb{C}\mathbb{P}^{m-1}$ coincides with f . This space is homotopy equivalent to the $2m$ -fold loop space of X and in what follows will be denoted by $\Omega^{2m} X$. We shall show that the inclusion

$$P_f \rightarrow \Omega^{2m} X$$

induces isomorphism in homology groups up to some dimension depending on m , X and the degrees of the polynomials contained in P_f . The precise statement of our theorem will be given in the next section.

In the case $m = 1$ our result includes the homology part of the theorem of Guest (for smooth varieties only) as a particular case and, moreover, gives an improvement over the results of [11] in what concerns the stabilization dimension.

2. Toric Varieties

A toric variety X is a complex algebraic variety which admits an action of the algebraic torus $(\mathbb{C}^*)^n$, with an open and dense orbit on which the action is free.

Given such X , we shall use the following notations:

- $N \simeq \mathbb{Z}^n$ — a lattice in \mathbb{R}^n .
- M — the integer dual of N .
- $\langle \cdot, \cdot \rangle : N \otimes M \rightarrow \mathbb{Z}$ — the natural pairing.
- $\Sigma \subset N_{\mathbb{R}} = \mathbb{R}^n$ — the fan of X .
- ρ_1, \dots, ρ_r — the one-dimensional cones in Σ .
- \mathbf{n}_i — the primitive generator of $\rho_i \cap N$.
- D_i — the divisor on X corresponding to ρ_i . (Recall that the 1-dimensional cones of Σ are in correspondence with irreducible $(\mathbb{C}^*)^n$ -invariant Weil divisors on X .)
- $A_{n-1}(X)$ — the group of Weil divisors modulo the subgroup of principal divisors. Two divisors $\sum a_i D_i$ and $\sum b_i D_i$ are in the same class in $A_{n-1}(X)$ if and only if there exists $\mathbf{m} \in M$ such that $a_i = \langle \mathbf{n}_i, \mathbf{m} \rangle + b_i$ for $i = 1, \dots, r$.

The details of the theory of toric varieties can be found in [5, 6, 7, 8].

In what follows we shall make extensive use of the homogeneous coordinate ring of X . It is defined as

$$R = \mathbb{C}[x_1, \dots, x_r],$$

where the variable x_i corresponds to ρ_i . Each monomial $\prod_i x_i^{a_i}$ determines a divisor

$$D = \sum_i a_i D_i;$$

we shall use the notation

$$x^D := \prod_i x_i^{a_i}.$$

We say that the monomial $\prod_i x_i^{a_i}$ has degree $[D] \in A_{n-1}(X)$; it follows that two monomials $\prod_i x_i^{a_i}$ and $\prod_i x_i^{b_i}$ have the same degree if and only if there is some $m \in M$ such that $a_i = \langle \mathbf{n}_i, m \rangle + b_i$.

In what follows we shall assume that X is a compact and smooth toric variety. Each cone $\sigma \in \Sigma$ determines the monomial $x^\sigma = \prod_{\rho_i \notin \sigma} x_i$. Define the subvariety

$$Y = \{(x_1, \dots, x_r) : x^\sigma = 0 \text{ for all } \sigma \in \Sigma\} \subset \mathbb{C}^r,$$

and the group

$$G = \left\{ (\mu_1, \dots, \mu_r) \in (\mathbb{C}^*)^r : \prod_{i=1}^r \mu_i^{\langle \mathbf{n}_i, m \rangle} = 1 \text{ for all } m \in M \right\}.$$

The group G acts on $\mathbb{C}^r - Y$ by multiplication coordinatewise. The quotient

$$X = (\mathbb{C}^r - Y)/G$$

is the toric variety associated to the fan Σ . This description gives us the homogeneous coordinates on X . In these coordinates the morphisms $\mathbb{C}\mathbb{P}^m \rightarrow X$ can be described explicitly as follows (see [5]).

Let $P_i : \mathbb{C}^{m+1} \rightarrow \mathbb{C}$ homogeneous polynomials for $i = 1, \dots, r$ such that P_i has degree d_i with $\sum_i d_i \mathbf{n}_i = 0$, and $(P_1(x), \dots, P_r(x)) \notin Y$ for all $x \in \mathbb{C}^{m+1} \setminus \{0\}$. Then the r -tuple (P_1, \dots, P_r) induces a morphism $\mathbb{C}\mathbb{P}^m \rightarrow X$. Furthermore,

- two sets of polynomials $\{P_i\}$ and $\{P'_i\}$ determine the same morphism if and only if there is $g \in G$ such that $(P_i) = g \cdot (P'_i)$;
- all morphisms $\mathbb{C}\mathbb{P}^m \rightarrow X$ arise in this way.

We say that a morphism $\mathbb{C}\mathbb{P}^m \rightarrow X$ has degree $\bar{d} = (d_1, \dots, d_r)$ if it is given by an r -tuple of homogeneous polynomials (P_i) with $\deg P_i = d_i$. Let $P_f(\bar{d})$ be the set of all morphisms $\mathbb{C}\mathbb{P}^m \rightarrow X$ of degree \bar{d} which restrict to a given morphism f on $\mathbb{C}\mathbb{P}^{m-1}$.

A set of edge generators $\{\mathbf{n}_{i_1}, \dots, \mathbf{n}_{i_j}\}$ for the fan Σ is *primitive* if it does not lie in any cone of Σ but every proper subset does. It can be proved that

$$Y = \bigcup_{\{\mathbf{n}_{i_1}, \dots, \mathbf{n}_{i_j}\} \text{ primitive}} \{(x_1, \dots, x_r) : x_{i_1} = \dots = x_{i_j} = 0\}.$$

We can now state the main result of the present paper.

Theorem 2.1. *Let X be a smooth compact toric variety associated to a fan Σ , with k the cardinality of the smallest primitive set of edge generators for Σ . Then, for all $m < k$, the inclusion of $P_f(\bar{d})$ into the corresponding component of $\Omega^{2m} X$ induces isomorphisms in homology groups for all dimensions smaller than $d(2k - 2m - 1) - 1$, where $\bar{d} = (d_1, \dots, d_r)$ and $d = \min\{d_i\}$.*

In the case of curves, that is, $m = 1$, our estimate is, in general, much stronger than the main theorem of Guest [11] and the result of Boyer, Hurtubise and Milgram [2] specialized to the case of toric varieties, both of which give the stabilization dimension d . Note, however, that the result of Guest also applies to singular toric varieties, while in our proof of Theorem 2.1 smoothness is essential: in the singular case we cannot, in general, represent morphisms by collections of polynomials.

The strategy of the proof is exactly the same as in [18] and [19] and mirrors other applications of Vassiliev’s method. It contains no essential novelty as compared to the case of a projective space [19], and thus we make an emphasis on the details specific to the case of a general toric variety.

First, we construct a sequence of finite-dimensional approximations to the space of all continuous maps from $\mathbb{C}\mathbb{P}^n$ to a toric variety. These finite-dimensional approximations are spaces of maps that are given by polynomials in both holomorphic and antiholomorphic variables, of fixed degrees, and the first in the sequence is the space of holomorphic maps. The Stone-Weierstrass Theorem implies that these approximations indeed have the space of all continuous maps as a topological limit.

The second part of the proof consists in comparing the topology of two successive finite-dimensional approximations. Here we make use of the fact that these spaces are complements of discriminants in affine spaces. In particular, their topology can be related to the topology of the corresponding discriminants by the Alexander duality. In order to calculate the cohomology of these discriminants one resolves their singularities with the help of the Vassiliev’s simplicial resolution, and considers the natural filtrations by the “fibrewise skeleta”.

The successive quotients of these filtrations are easy to describe and compare explicitly for the first few terms. For higher terms, it suffices to give a dimension estimate which allows to discard them.

In the next section we construct the spaces of finite-dimensional approximations and show that the limit space is, indeed, homotopy equivalent to the space of all continuous maps. Then, in Section 4 we describe the approximation spaces as complements of discriminants.

3. The Stone-Weierstrass Theorem

A (p, q) -polynomial in the variables z_i and \bar{z}_i (where $0 \leq i \leq m$) is a complex-valued homogeneous polynomial of degree p in the “holomorphic” variables z_i and degree q in “antiholomorphic” variables \bar{z}_i . If $\bar{p} = (p_1, \dots, p_r)$, $\bar{q} = (q_1, \dots, q_r) \in \mathbb{Z}^r$ with

$$\sum p_i \mathbf{n}_i = \sum q_i \mathbf{n}_i = 0,$$

we define a (\bar{p}, \bar{q}) -map as a map $F : \mathbb{C}\mathbb{P}^m \rightarrow X$ given by a r -tuple of (p_i, q_i) -polynomials such that $(F_1(x), \dots, F_r(x)) \notin Y$ for all $x \in \mathbb{C}^{m+1} \setminus \{0\}$.

Note that the $(\bar{d}, 0)$ -maps are the morphisms of degree \bar{d} . Two collections (P_i) and (P'_i) of (p_i, q_i) -polynomials determine the same morphism $F : \mathbb{C}\mathbb{P}^m \rightarrow X$ if and only if there are functions

$$g_1, \dots, g_r : \mathbb{C}^{m+1} \setminus \{0\} \rightarrow \mathbb{C}$$

such that $P_i = g_i P'_i$ and $(g_1(x), \dots, g_r(x)) \in G$ for all $x \in \mathbb{C}^{m+1} \setminus \{0\}$. In particular, the g'_i s do not vanish. Given

$$\bar{a} = (a_1, \dots, a_r) \in \mathbb{Z}^r \tag{1}$$

with

$$\sum_i a_i \mathbf{n}_i = 0,$$

a (\bar{p}, \bar{q}) -map can be written as a $(\bar{p} + \bar{a}, \bar{q} + \bar{a})$ -map by multiplying all the polynomials by $(|z|^{2a_1}, \dots, |z|^{2a_r})$ coordinatewise.

Let $f : \mathbb{C}\mathbb{P}^{m-1} \rightarrow X$ be a (\bar{p}, \bar{q}) -map and assume that we have chosen the (p_i, q_i) -polynomials f_i that define it. Denote by W_{p_i, q_i}^i the complex affine space of all (p_i, q_i) -polynomials whose restriction to $\mathbb{C}\mathbb{P}^{m-1}$ coincides with f_i , and by $W_{\bar{p}, \bar{q}}$ the cartesian product of the W_{p_i, q_i}^i . Let $P_f(\bar{p}, \bar{q}) \subset W_{\bar{p}, \bar{q}}$ be the subspace of all the r -tuples in $W_{\bar{p}, \bar{q}}$ whose values are not in Y for all points in $\mathbb{C}^{m+1} \setminus \{0\}$. These are precisely the r -tuples that define (\bar{p}, \bar{q}) -maps, though we stress that more than one r -tuple in $P_f(\bar{p}, \bar{q})$ may correspond to the same map.

For \bar{a} as in (1), we define the stabilization maps

$$\begin{aligned} P_f(\bar{p}, \bar{q}) &\rightarrow P_f(\bar{p} + \bar{a}, \bar{q} + \bar{a}), \\ (h_1(z), \dots, h_r(z)) &\mapsto (|z|^{2a_1} h_1(z), \dots, |z|^{2a_r} h_r(z)). \end{aligned}$$

The space $P_f(\bar{d} + \infty, \infty)$ is defined as the direct limit of these inclusions, where $\bar{d} = \bar{p} - \bar{q}$.

In the next section we shall see that these maps induce isomorphisms in homology in the dimensions given in Theorem 2.1. Here, we shall prove that the spaces $P_f(\bar{p}, \bar{q})$ approximate the space of all continuous maps topologically.

Proposition 3.1. *$P_f(\bar{d} + \infty, \infty)$ is homotopy equivalent to a connected component of $\Omega^{2m}(X)$.*

The principal tool in the proof of this statement is the Stone-Weierstrass Theorem for vector bundles (see, for instance, [18]).

Theorem 3.2. *Let E be a locally trivial real vector bundle over a compact space B , $s_\alpha : E \rightarrow B$ a set of its sections, and let A be a subalgebra of the \mathbb{R} -algebra $C(B)$ of continuous real-valued functions on B . Suppose that*

- the subalgebra A separates points of B , that is, for any pair $x, y \in B$ there exists $h \in A$ such that $h(x) \neq h(y)$,
- for any $y \in B$ there exists $h \in A$ such that $h(y) \neq 0$,
- for any $y \in B$ the fibre of E over y is spanned by $s_\alpha(y)$.

Then the A -module generated by the s_α is dense in the space of all continuous sections of E .

The Stone-Weierstrass Theorem has the following consequence.

Lemma 3.3. *Let W be a finite CW-complex. Any continuous map*

$$F : \mathbb{C}\mathbb{P}^m \times W \rightarrow X$$

can be uniformly approximated by a (p, q) -map, for some p and q , whose coefficients are functions on W . Moreover, if the restriction of F to $\mathbb{C}\mathbb{P}^{m-1} \times \{x\}$ is represented by a collection of polynomials $f_{i,x}$, the approximating polynomials can be chosen so as to coincide with $f_{i,x} \cdot |z|^{2k}$, for some k , on $\mathbb{C}\mathbb{P}^{m-1} \times W$.

The proof is entirely analogous to the proof in [18] for the case $X = \mathbb{C}\mathbb{P}^n$. Indeed, a continuous map $F : \mathbb{C}\mathbb{P}^m \times W \rightarrow X$ can be thought as a family of maps $\mathbb{C}\mathbb{P}^m \rightarrow X$ depending on a parameter $w \in W$. In particular, it can be given as a collection of $m + 1$ sections s_i of the line bundles $\mathcal{O}(d_i)$, with $\sum_{\rho} d_i \langle \mathbf{n}_i, \mathbf{m} \rangle = 0$ for some $\mathbf{m} \in M$, and each s_i depending on $w \in W$. Exactly as in [18], by the Stone-Weierstrass Theorem these sections can be approximated by functions, which, for any given w , are $(d_i + q, q)$ -polynomials for some q and these approximations can be chosen so as to give an arbitrarily close approximation of F .

Proof of Proposition 3.1. For any compact Riemannian manifold M and any space E there exists $\epsilon > 0$ such that any two maps $E \rightarrow M$ that are uniformly ϵ -close, are homotopic. Thus, Lemma 3.3 implies that the map of the homotopy groups

$$\pi_k P_f(\bar{d} + \infty, \infty) \rightarrow \pi_k \Omega^{2m}(X)$$

is surjective for all k . Setting $W = S^k$ we get that any element of $\pi_k \Omega^{2m}(X)$ can be approximated by a class in $\pi_k P_f(\bar{d} + \infty, \infty)$. On the other hand, this isomorphism is also injective, as any homotopy that goes through continuous maps can be approximated by a homotopy through (p, q) -maps (set $W = S^k \times [0, 1]$). Both spaces of maps have homotopy types of CW-complexes, so by the Whitehead Theorem they are homotopy equivalent. \square

4. The Space $P_f(\bar{p}, \bar{q})$ as a Complement to a Discriminant

In what follows the notation \mathbb{C}^m will be used for the affine chart $z_m = 1$ in $\mathbb{C}\mathbb{P}^m$. We shall use the notation T^\bullet for the one-point compactification of a space T .

Recall that we consider the toric variety X as the quotient $(\mathbb{C}^r - Y)/G$. The space $P_f(\bar{p}, \bar{q})$ is the complement of a discriminant Θ in the space $W_{\bar{p}, \bar{q}}$ which consists of collections of complex polynomials of multidegree \bar{p} in the variables z_i and \bar{q} in the \bar{z}_i .

$$\Theta = \Theta_{\bar{p}, \bar{q}} = \{(F_1, \dots, F_r) \in W_{\bar{p}, \bar{q}} : (F_1(z), \dots, F_r(z)) \in Y \text{ for some } z \in \mathbb{C}^m\}.$$

The cohomology groups of $P_f(\bar{p}, \bar{q})$ are related to the homology groups of the one-point compactification of Θ by the Alexander duality:

$$\tilde{H}^l(P_f(\bar{p}, \bar{q}), \mathbb{Z}) \simeq \tilde{H}_{2N_{\bar{p}, \bar{q}} - l - 1}(\Theta^\bullet, \mathbb{Z})$$

where $N_{\bar{p}, \bar{q}}$ is the complex dimension of $W_{\bar{p}, \bar{q}}$. A filtration Z_l^\bullet on Θ^\bullet gives rise to a spectral sequence converging to the homology of Θ^\bullet . In particular, the spectral sequence with the first term

$$E_{-r, s}^1 = \tilde{H}_{2N_{\bar{p}, \bar{q}} + r - s - 1}(\hat{Z}_r^\bullet / \hat{Z}_{r-1}^\bullet) \quad (2)$$

converges to the cohomology of $P_f(\bar{p}, \bar{q})$.

In order to construct a suitable filtration on Θ^\bullet , we shall use Vassiliev's method of simplicial resolutions [21]. Since not all the terms of the resulting filtration can be effectively described, we shall then use the truncation procedure as in [19].

Let $Z(\bar{p}, \bar{q}) = Z \subset W_{\bar{p}, \bar{q}} \times \mathbb{C}^m$ be the set

$$Z = \{(F_1, \dots, F_r, x) : (F_1(x), \dots, F_r(x)) \in Y\}.$$

There is a projection map $Z \rightarrow \Theta$ which forgets the point $x \in \mathbb{C}^m$. We denote by $Z^\Delta(\bar{p}, \bar{q})$, or simply by Z^Δ , the space of the non-degenerate simplicial resolution associated to this map. It is defined as the union of the spaces $Z_l = Z_l(\bar{p}, \bar{q})$, with positive l , which are constructed as follows. First, embed (non-linearly) the space \mathbb{C}^m into some vector space V in such a way that the image of any $2l$ distinct points of \mathbb{C}^m are not contained in any $2l - 2$ dimensional subspace. This gives an embedding

$$Z \hookrightarrow W_{\bar{p}, \bar{q}} \times \mathbb{C}^m \hookrightarrow W_{\bar{p}, \bar{q}} \times V.$$

The subspace $Z_l \subset W_{\bar{p}, \bar{q}} \times V$ consists of all the $l - 1$ -simplices which lie in the fibres of the projection map onto $W_{\bar{p}, \bar{q}}$ and whose vertices are on Z . We have a natural projection map

$$Z_l \rightarrow \Theta \subset W_{\bar{p}, \bar{q}}$$

whose fibres are $(l - 1)$ -skeleta of simplices of various, not necessarily finite, dimensions.

Consider the space of configurations of l distinct points in \mathbb{C}^m with labels in Y :

$$R_l := \{(x_i, s_i) \in (\mathbb{C}^m \times Y)^l : x_i \neq x_j \text{ for } i \neq j\} / S_l$$

where the symmetric group S_l acts by permuting the l coordinates $(x_1, s_1), \dots, (x_l, s_l)$. We write a point of R_l as $((x_i, s_i))$. Recall that Y can be seen as the union

$$Y = \bigcup_{\{\mathbf{n}_{i_1}, \dots, \mathbf{n}_{i_j}\} \text{ primitive}} \{(x_1, \dots, x_r) : x_{i_1} = \dots = x_{i_j} = 0\},$$

where $\mathbf{n}_1, \dots, \mathbf{n}_r$ are generators for the 1-dimensional cones of the fan Σ associated to X . Let k be the cardinality of the smallest primitive set of edge generators of Σ . Then R_l is a cell complex of dimension $2l(m + r - k)$.

Write p for $\min\{p_1, \dots, p_r\}$. We have the following

Proposition 4.1. *For $l \leq p$ the space $Z_l \setminus Z_{l-1}$ is an affine bundle over the space R_l , of real rank*

$$2(N_{\bar{p}, \bar{q}} - rl) + l - 1.$$

Proof. A point in

$$Z_l \setminus Z_{l-1} \subset W_{\bar{p}, \bar{q}} \times V$$

can be written as (F, t) where $F = (F_1, \dots, F_r)$ and t is in the convex hull in V of l distinct points $x_1, \dots, x_l \in \mathbb{C}^m$ with $F(x_j) \in Y$. There is a projection

$$\begin{aligned} Z_l \setminus Z_{l-1} &\rightarrow R_l \\ (F, t) &\mapsto ((x_j, F(x_j))). \end{aligned}$$

The inverse image of a fixed point $((x_j, s_j)) \in R_l$ is the cartesian product of the space of r -tuples of (p_i, q_i) -polynomials with

$$(F_1(x_j), \dots, F_r(x_j)) = s_j$$

and the convex hull of x_1, \dots, x_l in V . Each equation $F_i(x_j) = (s_j)_i$, with x_j and s_j fixed, is a linear equation in the space of coefficients W_{p_i, q_i} . If $l \leq p$, all these equations are affinely independent and, hence, the space of r -tuples of (p_i, q_i) -polynomials satisfying $F(x_j) = s_j$ has real dimension $2(N_{\bar{p}, \bar{q}} - rl)$. The convex hull of the x_j is an $l - 1$ simplex; adding the dimensions we get the proposition. \square

The stabilization map that sends $P_f(\bar{p}, \bar{q})$ to $P_f(\bar{p} + \bar{a}, \bar{q} + \bar{a})$ is, actually, the restriction of a map $W_{\bar{p}, \bar{q}} \rightarrow W_{\bar{p} + \bar{a}, \bar{q} + \bar{a}}$ which also sends $\Theta_{\bar{p}, \bar{q}}$ to $\Theta_{\bar{p} + \bar{a}, \bar{q} + \bar{a}}$.

This map also gives a map of the corresponding simplicial resolutions of the discriminants which respects the filtrations. It is clear from the proof of Proposition 4.1 that the structure of $Z_l \setminus Z_{l-1}$ as a bundle over R_l is preserved by the stabilization map. In particular, for $l \leq p$ the space

$$Z_l(\bar{p} + \bar{a}, \bar{q} + \bar{a}) \setminus Z_{l-1}(\bar{p} + \bar{a}, \bar{q} + \bar{a})$$

is an orientable affine bundle over

$$Z_l(\bar{p}, \bar{q}) \setminus Z_{l-1}(\bar{p}, \bar{q})$$

of real rank $2(N_{\bar{p}+\bar{a}, \bar{q}+\bar{a}} - N_{\bar{p}, \bar{q}})$. Since

$$(Z_l \setminus Z_{l-1})^\bullet = Z_l^\bullet / Z_{l-1}^\bullet,$$

we have that

$$\begin{aligned} \tilde{H}_i(Z_l(\bar{p}, \bar{q})^\bullet / Z_{l-1}(\bar{p}, \bar{q})^\bullet) = \\ \tilde{H}_{i+2(N_{\bar{p}+\bar{a}, \bar{q}+\bar{a}} - N_{\bar{p}, \bar{q}})}(Z_l(\bar{p} + \bar{a}, \bar{q} + \bar{a})^\bullet / Z_{l-1}(\bar{p} + \bar{a}, \bar{q} + \bar{a})^\bullet) \end{aligned} \quad (3)$$

for all i and $l \leq p$.

Consider the spectral sequences (2) for $P_f(\bar{p}, \bar{q})$ and for $P_f(\bar{p} + \bar{a}, \bar{q} + \bar{a})$. The isomorphism (3) implies that their entries coincide for all s and for $r \leq p$. There is little control over the terms of these spectral sequences when $r > p$. One way to correct for it is to replace the simplicial resolution of Θ by a truncated simplicial resolution as in [19]. Essentially, the truncated, after the p -th term, simplicial resolution consists in collapsing homotopically the non-contractible fibres of the natural projection

$$Z_p(\bar{p}, \bar{q})^\bullet \rightarrow \Theta(\bar{p}, \bar{q})^\bullet,$$

by means of adding a cone on each non-contractible fibre, see [19]. The truncated simplicial resolution has a filtration of length $p + 1$. Its first p terms coincide with those of the usual simplicial resolution and, according to Lemma 2.1 of [19], the last term is obtained from $Z_p(\bar{p}, \bar{q})^\bullet$ by adding cells of dimension at most

$$\dim(Z_p(\bar{p}, \bar{q}) \setminus Z_{p-1}(\bar{p}, \bar{q})) + 1.$$

By Proposition 4.1, this dimension equals

$$(2(N_{\bar{p}, \bar{q}} - pr) + p - 1) + \dim R_p + 1 = 2N_{\bar{p}, \bar{q}} + p(2m - 2k + 1).$$

Similarly, the corresponding truncated simplicial resolution for $\Theta(\bar{p} + \bar{a}, \bar{q} + \bar{a})^\bullet$ can be obtained by adding cells of dimension at most

$$2N_{\bar{p}+\bar{a}, \bar{q}+\bar{a}} + p(2m - 2k + 1)$$

to $Z_p(\bar{p}+\bar{a}, \bar{q}+\bar{a})^\bullet$. The effect of truncating the simplicial resolution on the spectral sequence is that all the terms $E_{-r,s}^1$ with $r > p + 1$ vanish, and $E_{-(p+1),s}^1$ can be non-zero only if $s \geq 2(k - m)p$. It follows from the definition of the Alexander duality pairing as the linking number that the map of spectral sequences induced by the stabilization map is an isomorphism for all $r \leq p$. As a result, we have

$$H^i(P_f(\bar{p}, \bar{q}), \mathbb{Z}) = H^i(P_f(\bar{p} + \bar{a}, \bar{q} + \bar{a}), \mathbb{Z})$$

for all $i < p(2k - 2m - 1) - 1$. If a map induces isomorphisms in integral cohomology up to some dimension, it also induces isomorphisms in integral homology in the same dimensions. We get

Proposition 4.2. *The homomorphism*

$$\tilde{H}_i(P_f(\bar{p}, \bar{q})) \rightarrow \tilde{H}_i(P_f(\bar{p} + \bar{a}, \bar{q} + \bar{a}))$$

induced by the stabilization map is an isomorphism for all $i < p(2k - 2m - 1) - 1$.

Since the direct limit of the stabilization maps is homotopy equivalent to the space of continuous maps by Proposition 3.1, Theorem 2.1 now follows.

Remark 4.3. In the case of curves, that is, $m = 1$, the stabilization dimension may be slightly improved. In this situation the space of based holomorphic curves is the complement to a discriminant which has a simplicial resolution with the filtration of length d , and the truncation procedure is not necessary. This argument, however, requires a different stabilization map, similar to that of [20]; we shall not discuss it here.

Remark 4.4. Spaces of based maps in Theorem 2.1 can be replaced by the corresponding spaces of free maps, with the same stabilization dimension; the argument is the same as that of Section 5 of [18].

Acknowledgements. This work is part of the doctoral thesis of the second author who was supported by a CONACyT scholarship. The first author was partially supported by the CONACyT grant 168093-F.

References

- [1] M. Adamaszek, A.Kozłowski, and K. Yamaguchi, *Spaces of Algebraic and Continuous Maps between Real Algebraic Varieties*, Quart. J. Math. **62** (2011), 771–790.
- [2] C. Boyer, J. Hurtubise, and J. Milgram, *Stability Theorems for Spaces of Rational Curves*, Internat. J. Math. **12** (2001), 223–262.

- [3] R. Cohen, J. D. S. Jones, and G. Segal, *Stability for Holomorphic Spheres and Morse Theory*, Contemp. Math. **258** (2000), 87–106.
- [4] R. Cohen, E. Lupercio, and G. Segal, *Holomorphic Spheres in Loop Groups and Bott Periodicity*, Asian J. Math. **3** (1999), 801–818.
- [5] D. A. Cox, *The Functor of a Smooth Toric Variety*, Tohoku Math. J. **47** (1995), 251–262.
- [6] ———, *The Homogeneous Coordinate Ring of a Toric Variety*, J. Algebraic Geom. **4** (1995), 17–50.
- [7] D. A. Cox, J. B. Little, and H. K. Schenck, *Toric Varieties*, Amer. Math. Soc, 2011.
- [8] W. Fulton, *Introduction to Toric Varieties*, Princeton University Press, 1993.
- [9] J. Gravesen, *On the Topology of Spaces of Holomorphic Maps*, Acta Math. **162** (1989), 247–286.
- [10] M. Guest, *Topology of the Space of Absolute Minima of the Energy Functional*, Amer. J. Math. **106** (1984), 21–42.
- [11] ———, *The Topology of the Space of Rational Curves on a Toric Variety*, Acta Math. **174** (1995), 119–145.
- [12] M. Guest, A. Kozłowski, and K. Yamaguchi, *Spaces of Polynomials with Roots of Bounded Multiplicity*, Fund. Math. **161** (1999), 93–117.
- [13] J. Hurtubise, *Holomorphic Maps of a Riemann Surface Into a Flag Manifold*, J. Diff. Geom. **43** (1996), 99–118.
- [14] F. Kirwan, *On Spaces of Maps From Riemann Surfaces to Grassmannians and Applications to the Cohomology of Moduli of Vector Bundles*, Ark. Mat. **24** (1986), 221–275.
- [15] B. Mann and J. Milgram, *Some Spaces of Holomorphic Maps to Complex Grassmann Manifolds*, J. Diff. Geom. **33** (1991), 301–324.
- [16] ———, *The Topology of Rational Maps to Grassmannians and a Homotopy Theoretic Proof of the Kirwan Stability Theorem*, Contemp. Math. **146** (1993), 251–275.
- [17] J. Mostovoy, *Spaces of Rational Loops on a Real Projective Space*, Trans. Amer. Math. Soc. **353** (2001), 1959–1970.
- [18] ———, *Spaces of Rational Maps and the Stone-Weierstrass Theorem*, Topology **45** (2006), 281–293.

- [19] ———, *Truncated Simplicial Resolutions and Spaces of Rational Maps*, Quart. J. Math. **63** (2012), 181–187.
- [20] G. Segal, *The Topology of Spaces of Rational Functions*, Acta Math. **143** (1979), 39–72.
- [21] V. A. Vassiliev, *Complements of Discriminants of Smooth Maps: Topology and Applications*, Translations of Mathematical Monographs, vol. 98, Amer. Math. Soc., 1992.

(Recibido en marzo de 2013. Aceptado en agosto de 2013)

DEPARTAMENTO DE MATEMÁTICAS
 CINVESTAV-IPN
 APARTADO POSTAL 14-740
 C.P. 07000, MEXICO
e-mail: jacob@math.cinvestav.mx

INSTITUT DE MATHÉMATIQUES DE JUSSIEU
 4, PLACE JUSSIEU, CASE 247
 75252 PARIS CÉDEX 5, FRANCE
e-mail: erendira.munguia@gmail.com

Esta página aparece intencionalmente en blanco